

1. 2008 AB2 “Concert ticket problem” [calculator-friendly]

t (hours)	0	1	3	4	7	8	9
$L(t)$ (people)	120	156	176	126	150	80	0

Concert tickets went on sale at noon ($t = 0$) and were sold out within 9 hours. The number of people waiting in line to purchase tickets at time t is modeled by a twice-differentiable function L for $0 \leq t \leq 9$. Values of $L(t)$ at various times t are shown in the table above.


- Use the data in the table to estimate the rate at which the number of people waiting in line was changing at 5:30 P.M. ($t = 5.5$). Show the computations that lead to your answer. Indicate units of measure.
- Use a trapezoidal sum with three subintervals to estimate the average number of people waiting in line during the first 4 hours that tickets were on sale.
- For $0 \leq t \leq 9$, what is the fewest number of times at which $L'(t)$ must equal 0? Give a reason for your answer.

2. 2008 AB4B [non-calculator]

The functions f and g are given by $f(x) = \int_0^{3x} \sqrt{4+t^2} dt$ and $g(x) = f(\sin x)$.

- Find $f'(x)$ and $g'(x)$.
- Write an equation for the line tangent to the graph of $y = g(x)$ at $x = \pi$.
- Write, but do not evaluate, an integral expression that represents the maximum value of g on the interval $0 \leq x \leq \pi$. Justify your answer.


3. 2005 AB5 [non-calculator]



A car is traveling on a straight road. For $0 \leq t \leq 24$ seconds, the car's velocity $v(t)$, in meters per second, is modeled by the piecewise-linear function defined by the graph above.

- Find $\int_0^{24} v(t) dt$. Using correct units, explain the meaning of $\int_0^{24} v(t) dt$.

4. 2008 AB5B [non-calculator]



Let g be a continuous function with $g(2) = 5$. The graph of the piecewise-linear function g' , the derivative of g , is shown above for $-3 \leq x \leq 7$.

- (a) Find the x -coordinate of all points of inflection of the graph of $y = g(x)$ for $-3 < x < 7$. Justify your answer.
- (b) Find the absolute maximum value of g on the interval $-3 \leq x \leq 7$. Justify your answer.
- (c) Find the average rate of change of $g(x)$ on the interval $-3 \leq x \leq 7$.
- (d) Find the average rate of change of $g'(x)$ on the interval $-3 \leq x \leq 7$. Does the Mean Value Theorem applied on the interval $-3 \leq x \leq 7$ guarantee a value of c , for $-3 < c < 7$, such that $g''(c)$ is equal to this average rate of change? Why or why not?

(e) If $f(x) = \int_{-1}^x g'(t) dt$, then what is the value of $f(2)$ [I added this part]

5. 2006 AB4 [non-calculator]

t (seconds)	0	10	20	30	40	50	60	70	80
$v(t)$ (feet per second)	5	14	22	29	35	40	44	47	49

Rocket A has positive velocity $v(t)$ after being launched upward from an initial height of 0 feet at time $t = 0$ seconds. The velocity of the rocket is recorded for selected values of t over the interval $0 \leq t \leq 80$ seconds, as shown in the table above.

- (a) Find the average acceleration of rocket A over the time interval $0 \leq t \leq 80$ seconds. Indicate units of measure.

- (b) Using correct units, explain the meaning of $\int_{10}^{70} v(t) dt$ in terms of the rocket's flight. Use a midpoint

Riemann sum with 3 subintervals of equal length to approximate $\int_{10}^{70} v(t) dt$.

6. 2007AB3 [calculator-friendly]

x	$f(x)$	$f'(x)$	$g(x)$	$g'(x)$
1	6	4	2	5
2	9	2	3	1
3	10	-4	4	2
4	-1	3	6	7

The functions f and g are differentiable for all real numbers, and g is strictly increasing. The table above gives values of the functions and their first derivatives at selected values of x . The function h is given by $h(x) = f(g(x)) - 6$.

- Explain why there must be a value r for $1 < r < 3$ such that $h(r) = -5$.
- Explain why there must be a value c for $1 < c < 3$ such that $h'(c) = -5$.
- Let w be the function given by $w(x) = \int_1^{g(x)} f(t) dt$. Find the value of $w'(3)$.

7. 2007 AB5 [non-calculator] In honor of Colorado's "balloon boy"

Let's do part (b) and (d) now


t (minutes)	0	2	5	7	11	12
$r'(t)$ (feet per minute)	5.7	4.0	2.0	1.2	0.6	0.5

The volume of a spherical hot air balloon expands as the air inside the balloon is heated. The radius of the balloon, in feet, is modeled by a twice-differentiable function r of time t , where t is measured in minutes. For $0 < t < 12$, the graph of r is concave down. The table above gives selected values of the rate of change, $r'(t)$, of the radius of the balloon over the time interval $0 \leq t \leq 12$. The radius of the balloon is 30 feet when $t = 5$.

(Note: The volume of a sphere of radius r is given by $V = \frac{4}{3}\pi r^3$.)

- Estimate the radius of the balloon when $t = 5.4$ using the tangent line approximation at $t = 5$. Is your estimate greater than or less than the true value? Give a reason for your answer.
- Find the rate of change of the volume of the balloon with respect to time when $t = 5$. Indicate units of measure.
- Use a right Riemann sum with the five subintervals indicated by the data in the table to approximate $\int_0^{12} r'(t) dt$. Using correct units, explain the meaning of $\int_0^{12} r'(t) dt$ in terms of the radius of the balloon.
- Is your approximation in part (c) greater than or less than $\int_0^{12} r'(t) dt$? Give a reason for your answer.

8. 2007 AB4 [non-calculator]




Let f be a function defined on the closed interval $-5 \leq x \leq 5$ with $f(1) = 3$. The graph of f' , the derivative of f , consists of two semicircles and two line segments, as shown above.

- For $-5 < x < 5$, find all values x at which f has a relative maximum. Justify your answer.
- For $-5 < x < 5$, find all values x at which the graph of f has a point of inflection. Justify your answer.
- Find all intervals on which the graph of f is concave up and also has positive slope. Explain your reasoning.
- Find the absolute minimum value of $f(x)$ over the closed interval $-5 \leq x \leq 5$. Explain your reasoning.

Hint for part (d): Don't forget the endpoints and consider how to find values of f with integration

9. 2006 AB3 "Periodic Trapezoid Problem" [non-calculator]



The graph of the function f shown above consists of six line segments. Let g be the function given by $g(x) = \int_0^x f(t) dt$.

- Find $g(4)$, $g'(4)$, and $g''(4)$.
- Does g have a relative minimum, a relative maximum, or neither at $x = 1$? Justify your answer.
- Suppose that f is defined for all real numbers x and is periodic with a period of length 5. The graph above shows two periods of f . Given that $g(5) = 2$, find $g(10)$ and write an equation for the line tangent to the graph of g at $x = 108$.

Some multiple-choice questions to practice:

5. 1985 AB42

$$\frac{d}{dx} \int_2^x \sqrt{1+t^2} dt =$$

(A) $\frac{x}{\sqrt{1+x^2}}$ (B) $\sqrt{1+x^2} - 5$ (C) $\sqrt{1+x^2}$ (D) $\frac{x}{\sqrt{1+x^2}} - \frac{1}{\sqrt{5}}$

(E) $\frac{1}{2\sqrt{1+x^2}} - \frac{1}{2\sqrt{5}}$

6. 1988 AB13

If the function f has a continuous derivative on $[0, c]$, then $\int_0^c f'(x) dx =$

(A) $f(c) - f(0)$ (B) $|f(c) - f(0)|$ (C) $f(c)$ (D) $f(x) + c$

(E) $f''(c) - f''(0)$

8. 1988 BC14

If $F(x) = \int_1^{x^2} \sqrt{1+t^3} dt$, then $F'(x) =$

(A) $2x\sqrt{1+x^6}$ (B) $2x\sqrt{1+x^3}$ (C) $\sqrt{1+x^6}$ (D) $\sqrt{1+x^3}$


(E) $\int_1^{x^2} \frac{3t^2}{2\sqrt{1+t^3}} dt$

9. 1993 AB41

$\frac{d}{dx} \int_0^x \cos(2\pi u) du$ is

(A) 0 (B) $\frac{1}{2\pi} \sin x$ (C) $\frac{1}{2\pi} \cos(2\pi x)$ (D) $\cos(2\pi x)$ (E) $2\pi \cos(2\pi x)$


11. 1997 AB78



The graph of f is shown in the figure above. If $\int_1^3 f(x) dx = 2.3$ and $F'(x) = f(x)$, then $F(3) - F(0) =$

- (A) 0.3 (B) 1.3 (C) 3.3 (D) 4.3 (E) 5.3

20. 2003 AB22



The graph of f' , the derivative of f , is the line shown in the figure above. If $f(0) = 5$, then $f(1) =$

- (A) 0 (B) 3 (C) 6 (D) 8 (E) 11